



ELSEVIER

# Effects of nucleoid-associated proteins on bacterial chromosome structure and gene expression

Douglas F Browning<sup>1</sup>, David C Grainger<sup>2</sup> and Stephen JW Busby<sup>3</sup>

Bacterial nucleoid-associated proteins play a key role in the organisation, replication, segregation, repair and expression of bacterial chromosomes. Here, we review some recent progress in our understanding of the effects of these proteins on DNA and their biological role, focussing mainly on *Escherichia coli* and its chromosome. Certain nucleoid-associated proteins also regulate transcription initiation at specific promoters, and work in concert with dedicated transcription factors to regulate gene expression in response to growth phase and environmental change. Some specific examples, involving the *E. coli* IHF and Fis proteins, that illustrate new principles, are described in detail.

## Addresses

<sup>1</sup> School of Immunity and Infection, University of Birmingham, Birmingham B15 2TT, UK

<sup>2</sup> Department of Biological Sciences, University of Warwick, Coventry CV4 7AL, UK

<sup>3</sup> School of Biosciences, University of Birmingham, Birmingham B15 2TT, UK

Corresponding author: Busby, Stephen JW ([s.j.w.busby@bham.ac.uk](mailto:s.j.w.busby@bham.ac.uk))

Current Opinion in Microbiology 2010, 13:773–780

This review comes from a themed issue on  
Growth and development: prokaryotes  
Edited by Joe Lutkenhaus

Available online 13th October 2010

1369-5274/\$ – see front matter

© 2010 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.mib.2010.09.013

## Introduction

The nucleoid is a region of the bacterial cell into which its chromosome is constrained. To fit into this space, the DNA has to be highly compacted and this is due to the action of supercoiling, RNA and nucleoid-associated proteins (NAPs). Bacterial chromosome folding and the factors involved in their folding began to be investigated in the 1970s. However, the first thorough and systematic investigation of NAPs came from Akira Ishihama and colleagues in 1999 [1,2], who catalogued the different NAPs of *Escherichia coli*, measured their binding affinities, investigated their preferred targets, quantified their levels, and reported dramatic changes in the levels of certain NAPs in response to changes in growth. The arrival of chromatin immunoprecipitation technology together with whole genome sequences has led to a picture of the distribution of individual NAPs across

whole chromosomes [3], whilst the total protein occupancy landscape has been established using complementary experimental approaches [4,5]. Remarkably, we are still largely ignorant of precisely how bacterial chromosomes are packaged and organised, but the literature, perhaps mistakenly, places a big emphasis on the importance of NAPs [6]. Here, we present a selection of recent reports, describing effects of NAPs on bacterial chromosome structure and gene expression, and we refer the reader to the excellent review by Dillon and Dorman [7] for a comprehensive up-to-date picture.

## Bacterial folded chromosomes

Of all the bacterial chromosomes, it is the *E. coli* chromosome that has been most studied. Its packaging and organisation can be probed using fluorescent proteins targeted to specific loci and Sherratt and colleagues' recent review [8] explains how its replication and segregation are organised. Wiggins *et al.* [9] have pushed the technology to the limit and measured the distribution and fluctuations of many different locations and their juxtaposition, and they conclude that loci show remarkable precision in their positioning. Thus, although many textbooks sketch bacterial chromosomes as a more or less disordered jumble of DNA, this is far from the reality and they are likely to be organised with the same precision as other macromolecular assemblies, although with a built-in facility for change due to the mechanics of DNA replication, repair, segregation and expression. With respect to building this structure, Sexton *et al.* [10] argue for the importance of meeting points between different locations in chromosomes, and Junier *et al.* [11] describe the physical consequences of these interactions. The resulting periodicity, and the creation of different neighbourhoods within the same cell, may have consequences for gene expression and for the organisation of regulatory networks, and these are explored by Janga *et al.* [12] and Mathelier and Carbone [13].

The traditional view of the *E. coli* folded chromosome is that it consists of a series of constrained loops separated by dynamic boundaries. Boccard and his colleagues have suggested a higher order of organisation in which the chromosome is partitioned into six discrete 'macrodomains', and fluorescence microscopy shows that the local DNA dynamics differs in each of the macrodomains that may correspond to distinct physical structures [14]. In a recent study [15], Boccard's team identified a new nucleoid-associated protein, MatP, whose binding is specific for the *ter* macrodomain that covers the replica-

tion termination region. This is an important result as it shows that the local structure of each segment of a bacterial chromosome might not be identical, though, for the moment, it is not clear how MatP binding confers distinctive properties on the *ter* macrodomain.

### Flexibility and variation in NAPs

Studies of different bacterial systems in different conditions have revealed complexities that were not apparent in Ishihama and colleagues' original 'audit' of *E. coli* NAPs [1,2]. For example, the observation that the high levels of Fis in rapidly growing cells are reduced to near zero as cell growth slows does not apply in anaerobic cultures of *Salmonella enterica* serovar Typhimurium [16]. Moreover, there is great variation in NAPs in different bacteria and Fis is restricted to Gamma proteobacteria [17], although AbrB appears to play the same role in *Bacillus subtilis* [18]. Takeyasu and colleagues have exploited atomic force microscopy [19] and optical tweezers [20] to observe big changes in the fibre structure of the *E. coli* chromosomes in different growth conditions and have attributed these to changes in the levels of different NAPs [17] and also transcription patterns [21,22]. However the behaviour of *E. coli* is more elaborate than many bacteria and, for example, such changes are not seen in the chromosomes of *Staphylococcus aureus* [23]. Finally the lists of different NAPs found in the literature for different bacteria may not yet be a closed book. Thus the *B. subtilis* Noc protein [24] and the *E. coli* Dan protein [25] are recent additions, and the discovery of MatP [15\*] hints that there may be many more locally acting NAPs to be found.

### NAPs and DNA structure

*In vitro* studies with purified individual NAPs using both ensemble and single molecule measurements show that bending, bridging, wrapping and clustering can result, following binding to DNA [6]. Some of these modes are illustrated in Figure 1. For example, DNA bridging by H-NS has been shown most comprehensively by Remus Dame and his colleagues using a variety of biophysical methods [26]. A recent report suggests that different modes of H-NS action could be controlled by divalent metal ions [27\*]. Similarly, HU, the most universally conserved and abundant NAP, can induce DNA bends, condense DNA in a fibre, and also interact with single stranded DNA [28,29]. Combinations of NAPs have been studied systematically by Muskhelishvili and colleagues and shown to generate novel types of DNA organisation [30]. Significantly, Dps and CbpA, both of which are strongly induced in stationary phase *E. coli*, cause compaction by clustering of distal DNA loci (Figure 1b) [31].

### NAPs and global transcription

Since most of the NAPs are high abundance and quite promiscuous with respect to DNA binding it is unsurprising that they affect transcription on a global scale. For


example, two recent reports [32,33] underscore the profound influence of HU in coordinating the *E. coli* transcriptome. The importance of NAPs in global transcription is reinforced by chromatin immunoprecipitation experiments which showed that, for H-NS, IHF and Fis in *E. coli*, the majority of binding targets were located in intergenic regulatory regions [34]. Effects on global transcription not only occur due to changes in the relative levels of NAPs in different growth phases, but also arise due to NAP binding being affected by the passage of replication forks during the cell cycle. For example, a recent report [35] highlighted a possible role for SeqA, which binds to targets containing hemi-methylated GATC motifs.

Much recent literature attention has focused on H-NS, which is often referred to as the 'genome guardian', the 'universal repressor' or a 'sentinel' [7\*,36]. H-NS presents a tantalising problem, because, despite having been first reported in 1971 [37], we still lack understanding, at the molecular level, of specific repression by H-NS and how it is modulated. What is clear is that H-NS recognizes AT-rich segments of the chromosome, and then binding is in two steps, an initial binding of the H-NS dimer, followed by further co-operative binding, and this can result in molecular bridging between distal chromosome locations and concomitant transcription regulation [36]. A major breakthrough was the identification of high affinity DNA sites for H-NS that are, perhaps, responsible for step one of H-NS binding [38\*].

The existence of the H-NS paralogue, StpA, in both *E. coli* and *S. enterica* serovar Typhimurium has long posed a puzzle, since *stpA* knockouts have minimal effects in *E. coli*. Using biochemical approaches, John Ladbury's group showed that H-NS and StpA could form both homodimers and heterodimers [39] and this may well explain genome-wide chromatin immunoprecipitation data from Oshima and colleagues [40], who found that the vast majority of DNA targets in *E. coli* occupied by StpA were also occupied by H-NS. Deletion of *stpA* had a minimal effect on the distribution of H-NS, whilst deletion of *hns* greatly reduced the number of StpA-bound targets. Thus, in *E. coli*, StpA binds mainly as a heterodimer with H-NS, acting almost as a 'reserve'. By contrast, in *S. enterica* serovar Typhimurium, StpA plays a significant role, an *stpA* deletion affects expression of 5% of the genome, and there appears to be an StpA regulon [41].

The observation that many of the A:T-rich targets for H-NS are located in genome segments that have arrived by horizontal evolution prompted the idea that H-NS is a 'sentinel' whose task is to suppress expression from externally acquired pieces of DNA [7\*]. Support for this comes from the observation that some transmissible plasmids carry a gene encoding an H-NS homologue. Doyle

Figure 1



DNA folding properties of bacterial nucleoid proteins. **(a)** Log phase-specific and constitutively expressed nucleoid proteins. The diagrams depict the DNA folding properties of nucleoid proteins (shown as grey spheres) found either in only growing cells (Fis) or in all phases of growth (IHF, HU and H-NS). **(b)** Stationary phase-specific nucleoid proteins. The figure depicts the DNA folding properties of the nucleoid proteins Dps and CbpA (shown as grey spheres). Both proteins self-aggregate when associated with DNA.

*et al.* [42<sup>\*</sup>] identified Sfh as such a homologue and showed that, without Sfh, incoming plasmids stress the host cell as they titrate out host H-NS. Hence Sfh can be viewed as a molecular stealth device that permits the plasmid to enter the host without causing disruption. A recent genome-wide analysis of H-NS and Sfh binding in *S. enterica* serovar Typhimurium [43] supports this view by showing that Sfh and H-NS recognise common target sites, and argues that the invading plasmid deploys Sfh to 'top-up' the level of H-NS thereby minimizing disturbance to its new host. This scenario, with H-NS as a sentinel, supposes that, if a segment of horizontally acquired DNA segment encoded something that was useful for its new host, mechanisms would evolve to ensure activation. Recent reviews from Dorman and colleagues [44,45] discuss this issue and outline the different mechanisms by which repression by H-NS and its homologues at specific targets can be lifted.

Another current focus of H-NS research concerns *Mycobacteria* where initial scans of genome sequences suggested that H-NS was absent and that a protein that resembled the GroEL chaperone functioned as a NAP [46]. However it has now been shown that the *M. tuberculosis* Rv3597 and Rv3852 genes encode proteins (Lsr2 and H-NS respectively) whose *in vitro* properties resemble H-NS from enterics. Concerning Lsr2, although it shares but 4% sequence identity with *E. coli* H-NS, it is a DNA-bridging protein that binds to AT-rich regions, and can complement some effects of *hns* mutations in *E. coli* [47,48]. Similarly, the H-NS-like product of the *M. tuberculosis* Rv3852 gene can complement *hns* mutations in *E. coli* [49]. Interestingly the *M. tuberculosis* Rv3852 gene product has been shown to interact with Holliday junctions [50], and this underscores the point that, although the majority of literature concerning H-NS and its paralogues focuses on their interaction with double


stranded DNA, there is considerable evidence for interactions with single stranded nucleic acids, including messenger RNA. For example, a recent report [51<sup>\*</sup>] describes interactions of H-NS with the leader sequence of the *E. coli maltT* gene, and argues that H-NS activates the initiation of translation of MalT, and may interact likewise at other messengers.

### IHF: a NAP that likes to play with others

Bacterial Integration Host Factor (IHF) is a NAP that plays many roles, binds at many DNA targets, and is known to act like a transcription factor at many gene regulatory regions in *E. coli* [34]. Its major function is to bend target DNA sharply and hence it often functions in conjunction with other NAPs and transcription factors, such as at the *fim* switch that controls phase variable of type 1 fimbriae in *E. coli* [52]. Recent experiments with the *E. coli nir* and *nrf* promoters, which both are induced by anaerobiosis together with nitrite or nitrate ions, have revealed new subtleties in IHF action. Both promoters control the expression of a nitrite reductase and expres-


sion is completely dependent on FNR, the master regulator of transcription in anaerobic conditions. At both promoters, FNR-dependent transcription activation is suppressed by upstream-bound IHF. At the *nir* promoter, this suppression is effected by the combined action of IHF (which binds at position  $-88$ ) and Fis (which binds at position  $-142$ ). The dependence of the promoter on nitrite or nitrate ions comes from the fact that NarL and NarP, transcription activators that are triggered by nitrite and nitrate, bind to a site that overlaps the DNA site for IHF (Figure 2a). Binding displaces IHF and this results in relief of suppression of FNR-dependent *nir* promoter activity. The unexpected subtlety is the existence of a second weaker DNA site for IHF located at position  $-115$ , and that IHF binding at this site is activatory for FNR-dependent *nir* promoter activity. Hence basal *nir* promoter activity in anaerobic conditions in the absence of nitrite or nitrate depends on the relative occupation of the two DNA sites for IHF. The base sequence of these two sites varies from one *E. coli* strain to another and this variation sets the basal activity in any strain [53].

Figure 2



Transcription regulation at the *E. coli nir* and *nrf* operon promoters. (a) The *nir* promoter. Transcription initiation is dependent on FNR. The binding of Fis and IHF inhibits FNR-activated transcription (-ve), whilst IHF binding to a second lower affinity site stimulates transcription (+ve). The binding of NarL (or its homologue, NarP) displaces IHF, counteracting the repression mediated by IHF and Fis, and enabling maximal FNR-dependent transcription. (b) The *nrf* promoter. Transcription initiation is dependent on FNR. The binding of IHF, Fis and NsrR to the promoter inhibits FNR-activated transcription (-ve), whilst occupancy of a second upstream IHF site stimulates transcription (+ve). The binding of NarL or NarP displaces IHF, counteracting IHF-mediated repression and increasing FNR-dependent transcription. Repression by NsrR is relieved by the presence of reactive nitrogen species. In both panels, the transcription start point (+1) is indicated by a bent arrow.

Figure 3



Regulation of transcription initiation by Fis. The figure shows sketches to illustrate the different ways that Fis can interact at bacterial promoters. In all panels, the transcription startpoint (+1) is indicated by a bent arrow. **(a)** Simple class I activation. Fis directly contacts the C-terminal domain of one of the  $\alpha$  subunits of holo RNA polymerase ( $E\sigma^{70}$ ), thereby recruiting RNA polymerase to the promoter. **(b)** Altering DNA conformation. Binding of Fis to the upstream of the promoter stabilizes the “breathing” of the DNA helix, transmitting this energy downstream and facilitating the ease with RNA polymerase interacts with the promoter. **(c)** Direct repression. The binding of Fis to the promoter region directly blocks the association of holo RNA polymerase ( $E\sigma^{70}$ ). **(d)** Displacement of an essential activator. Fis represses transcription initiation by displacing an essential activator (A). **(e)** Preventing the function of an activator. Fis represses transcription initiation by interfering with the ability of an activator protein (A) to activate transcription. **(f)** Jamming of RNA polymerase at promoters. Binding of Fis, in unison with holo RNA polymerase containing  $\sigma^{70}$  ( $E\sigma^{70}$ ), shuts down transcription by creating a repression complex. This blocks access to the promoter by RNA polymerase containing the alternative “stationary phase”  $\sigma^{38}$  factor ( $E\sigma^{38}$ ).

A similar situation is found at the *E. coli nrf* promoter, where IHF bound at position -54 represses FNR-dependent activation, whilst IHF bound further upstream at -127 promotes activation. Surprisingly, the repression of

FNR-dependent activation by IHF at the *nrf* promoter is less efficient than at the *nir* promoter, and it has recently been shown that a second repressor, NsrR is involved [54]. Hence, FNR-dependent activation of the *nrf* pro-



moter is down-regulated independently by two different factors, IHF that binds at position  $-54$ , and NsrR that binds at position  $-63$ . Repression by IHF is lifted by NarL or NarP binding in response to nitrate or nitrite, whilst repression by NsrR is lifted when its DNA binding activity is prevented when it interacts with reactive nitrogen species generated as by-products of nitrate or nitrite metabolism (Figure 2b). A distinctive feature of the *E. coli nrf* promoter is that binding of Fis to a target at position  $-15$  overrides all the other inputs from FNR, IHF, NsrR, NarL and NarP, and thus, unusually, this promoter is subject to three different methods of repression (Figure 2b).

### Fis: the busy-body of the cell

The factor for inversion stimulation (Fis) is another DNA-bending NAP that plays many roles and binds at hundreds of DNA targets in *E. coli* [34,55]. It is especially important because it reaches very high levels in rapidly growing cells but it is almost absent in certain conditions of slow growth. The first crystal structure of Fis bound to a DNA target has recently been reported [56<sup>\*</sup>]. Detailed studies of Fis action at individual regulatory regions have revealed at least six different mechanisms by which it affects gene expression (summarised in Figure 3). At some promoters, Fis activates transcription initiation either by directly interacting with RNA polymerase thereby recruiting it to the promoter, or by inducing an activatory conformational change in the promoter [57]. However a major task for Fis appears to be to shut off the expression of inessential gene products during rapid growth. This can be done by direct binding to essential promoter elements as at the *E. coli nrf* promoter (Figure 2b), or by displacing an essential activator [58] or preventing the function of an activator [53]. Another mechanism of repression employed by Fis is RNA polymerase jamming. Hence at the *E. coli dps* promoter, that controls expression of the major stationary phase NAP, Dps, Fis combines with RNA polymerase containing the house keeping sigma 70 factor to form a repressive ternary complex [59].

### Conclusions and perspectives

Although our understanding of NAPs and their effects has greatly advanced over the past few years, there are still many issues to be tackled. Notably, how is bacterial chromosomal DNA organised in cells and what are the consequences on replication, transcription and repair? What do the different macrodomains correspond to and how are they controlled? How far is the *E. coli* model applicable to other bacteria where there are a smaller number of NAP species and do other factors such as cis-acting RNA play a role? Many of these issues are dealt with in a stimulating way in a recently published edited book on bacterial chromatin [60]. Meanwhile many fresh approaches are being taken to tackle the problems, including sophisticated imaging, tomography and new ana-

lytical approaches to home in on selected segments of bacterial chromosomes [61].

### Acknowledgements

The authors have been supported by a Wellcome Trust Programme Grant to SJWB and a Wellcome Trust Research Career Development Fellowship to DCG. We apologise that we had to omit many excellent relevant papers due to space constraints.

### References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
1. Azam TA, Ishihama A: **Twelve species of the nucleoid-associated protein from *Escherichia coli*. Sequence recognition specificity and DNA binding affinity.** *J Biol Chem* 1999, **274**:33105-33113.
  2. Azam TA, Iwata A, Nishimura A, Ueda S, Ishihama A: **Growth phase-dependent variation in protein composition of the *Escherichia coli* nucleoid.** *J Bacteriol* 1999, **181**:6351-6370.
  3. Grainger DC, Busby SJ: **Global regulators of transcription in *Escherichia coli*: mechanisms of action and methods for study.** *Adv Appl Microbiol* 2008, **65**:93-113.
  4. Vora T, Hottes AK, Tavazoie S: **Protein occupancy landscape of a bacterial genome.** *Mol Cell* 2009, **35**:247-253.  
The authors use a novel approach to locate all the regions of the *E. coli* chromosome that are bound by protein. The study identifies extended loci with high protein occupancy that are transcriptionally silent. Ref. [5] gives an insightful commentary on this paper.
  5. Cho B-K, Palsson BO: **Can the protein occupancy landscape show the topologically isolated chromosomal domains in the *E. coli* genome? An exciting prospect.** *Mol Cell* 2009, **35**:255-256.
  6. Luijsterbeurg MS, White MF, van Driel R, Dame RT: **The major architects of chromatin: architectural proteins in bacteria, archaea and eukaryotes.** *Crit Rev Biochem Mol Biol* 2008, **43**:393-418.
  7. Dillon SC, Dorman CJ: **Bacterial nucleoid-associated proteins, nucleoid structure and gene expression.** *Nat Rev Microbiol* 2010, **8**:185-195.  
This easy to read, yet comprehensive review, details the many roles that NAPs play in structuring the bacterial chromosome and regulating gene expression in different bacterial species.
  8. Reyes-Lamothe R, Wang X, Sherratt D: ***Escherichia coli* and its chromosome.** *Trends Microbiol* 2008, **16**:238-245.
  9. Wiggins PA, Cheveralls KC, Martin JS, Lintner R, Kondev J: **Strong intranucleoid interactions organise the *Escherichia coli* chromosome into a nuclear filament.** *Proc Natl Acad Sci* 2010, **107**:4991-4995.
  10. Sexton T, Bantignies F, Cavalli G: **Genomic interactions: Chromatin loops and gene meeting points in transcriptional regulation.** *Semin Cell Dev Biol* 2009, **20**:849-855.
  11. Junier I, Martin O, Kepes F: **Spatial and topological organisation of DNA chains induced by gene co-localization.** *PLoS Comput Biol* 2010, **6**:e1000678.
  12. Janga SC, Selgado H, Martinez-Antonio A: **Transcriptional regulation shapes the organization of genes on bacterial chromosomes.** *Nucleic Acids Res* 2009, **37**:3680-3688.
  13. Mathelier A, Carbone A: **Chromosomal periodicity and positional networks of genes in *Escherichia coli*.** *Mol Syst Biol* 2010, **6**:366.
  14. Espeli O, Mercier R, Boccard F: **DNA dynamics vary according to macrodomain topography in the *E. coli* chromosome.** *Mol Microbiol* 2008, **68**:1418-1427.
  15. Mercier R, Petit MA, Schbath S, Robin S, El Karoul M, Boccard F, Espeli O: **The MatP/mats site-specific system organises the**

- terminus region of the *E. coli* chromosome into a macrodomain.** *Cell* 2008, **135**:475-485.
- This exciting paper describes the ingenious discovery of the MatP protein, which binds to targets specifically located in an extended region of the *E. coli* chromosome centred on the replication terminus.
16. Croinin TO, Dorman CJ: **Expression of the Fis protein is sustained in late-exponential and stationary-phase cultures of *Salmonella enterica* serovar Typhimurium grown in the absence of aeration.** *Mol Microbiol* 2007, **66**:237-251.
  17. Ohniwa RL, Morikawa K, Kim J, Ohta T, Ishihama A, Wada C, Takeyasu K: **Dynamic state of DNA topology is essential for genome condensation in bacteria.** *EMBO J* 2006, **25**:5591-5602.
  18. O'Reilly M, Devine KM: **Expression of AbrB, a transition state regulator from *Bacillus subtilis*, is growth phase dependent in a manner resembling that of Fis, the nucleoid binding protein from *Escherichia coli*.** *J Bacteriol* 1997, **179**:522-529.
  19. Kim J, Yoshimura SH, Hizume K, Ohniwa RL, Ishihama A, Takeyasu K: **Fundamental structural units of the *Escherichia coli* nucleoid revealed by atomic force microscopy.** *Nucleic Acids Res* 2004, **32**:1982-1992.
  20. Shindo E, Kubo K, Ohniwa RL, Takeyasu K, Yoshikawa K: **In situ analysis of the higher-order genome structure in a single *Escherichia coli* cell.** *J Biotechnol* 2008, **133**:172-176.
  21. Ohniwa RL, Morikawa K, Takeshita SL, Kim J, Ohta T, Wada C, Takeyasu K: **Transcription-coupled nucleoid architecture in bacteria.** *Genes Cells* 2007, **12**:1141-1152.
  22. Cabrera JE, Cagliero C, Quan S, Squires CL, Jin DJ: **Active transcription of rRNA operons condenses the nucleoid in *Escherichia coli*: examining the effect of transcription on nucleoid structure in the absence of transcription.** *J Bacteriol* 2009, **191**:4180-4185.
  23. Morikawa K, Ohniwa RL, Kim J, Maruyama A, Ohta T, Takeyasu K: **Bacterial nucleoid dynamics: oxidative stress response in *Staphylococcus aureus*.** *Genes Cells* 2006, **11**:409-423.
  24. Wu LJ, Ishikawa S, Kawai Y, Oshima T, Ogasawara N, Errington J: **Noc protein binds to specific DNA sequences to coordinate cell division with chromosome segregation.** *EMBO J* 2009, **28**:1940-1952.
  25. Teramoto J, Yoshimura SH, Takeyasu K, Ishihama A: **A novel nucleoid protein of *Escherichia coli* induced under anaerobic growth conditions.** *Nucleic Acids Res* 2010, **38**:3605-3618.
  26. Wiggins PA, Dame RT, Noom MC, Wuite GJ: **Protein-mediated molecular bridging: a key mechanism in biopolymer organization.** *Biophys J* 2009, **97**:1997-2003.
  27. Liu Y, Chen H, Kenney LJ, Yan J: **A divalent switch drives H-NS/DNA-binding conformations between stiffening and bridging modes.** *Genes Dev* 2010, **24**:339-344.
- It has long been known that the interaction between H-NS and its DNA targets can be influenced by salt and temperature. This paper argues that divalent metal ions may play an important role in switching H-NS between different binding modes.
28. van Noort J, Verbrugge S, Goosen N, Dekker C, Dame RT: **Dual architectural roles of HU: formation of flexible hinges and rigid filaments.** *Proc Natl Acad Sci* 2004, **101**:6969-6974.
  29. Kamashev D, Balandina A, Mazur AK, Arimondo PB, Rouviere-Yaniv J: **HU binds and folds single-stranded DNA.** *Nucleic Acids Res* 2008, **36**:1026-1036.
  30. Maurer S, Fritz J, Muskhelishvili G: **A systematic *in vitro* study of nucleoprotein complexes formed by bacterial nucleoid-associated proteins revealing novel types of DNA organization.** *J Mol Biol* 2009, **387**:1261-1276.
  31. Cosgriff S, Chintakayala K, Chim YTA, Chen X, Allen S, Lovering AL, Grainger DC: **Dimerisation and DNA-dependent aggregation of the *E. coli* nucleoid protein and chaperone CbpA.** *Mol Microbiol* 2010, **77**:1289-1300.
  32. Berger M, Farcas A, Geertz M, Zhelyazkova P, Brix K, Travers A, Muskhelishvili G: **Coordination of genomic structure and transcription by the main bacterial nucleoid-associated protein HU.** *EMBO Rep* 2009, **11**:59-64.
  33. Oberto J, Nabti S, Jooste V, Mignot H, Rouviere-Yaniv J: **The HU regulon is composed of genes responding to anaerobiosis, acid stress, high osmolarity and SOS induction.** *PLoS One* 2009, **34**:e4367.
  34. Grainger DC, Hurd D, Goldberg MD, Busby SJ: **Association of nucleoid proteins with coding and non-coding segments of the *Escherichia coli* genome.** *Nucleic Acids Res* 2006, **34**:4642-4652.
  35. Sanchez-Romero MA, Busby SJW, Dyer NP, Ott S, Millard AD, Grainger DC: **Dynamic distribution of SeqA protein across the chromosome of *Escherichia coli* K-12.** *mBio* 2010, **1**: e0012-10.
  36. Fang FC, Rimsky S: **New insights into transcriptional regulation by H-NS.** *Curr Opin Microbiol* 2008, **11**:113-120.
  37. Jacquet M, Cukier-Kahn R, Pla J, Gros F: **A thermostable factor acting on *in vitro* DNA transcription.** *Biochem Biophys Res Commun* 1971, **45**:1597-1607.
  38. Lang B, Blot N, Bouffartigues E, Buckle M, Geertz M, Gualerzi GO, Mavathur R, Muskhelishvili G, Pon C, Rimsky S *et al.*: **High-affinity DNA binding sites for H-NS provide a molecular basis for selective silencing within proteobacterial genomes.** *Nucleic Acids Res* 2007, **35**:6330-6337.
- In this clever application of bioinformatics, the authors build on their previous observation that H-NS binds to DNA by recognising a specific "nucleation" sequence.
39. Leonard PG, Ono S, Gor J, Perkins SJ, Ladbury JE: **Investigation of the self-association and hetero-association interactions of H-NS and StpA from Enterobacteria.** *Mol Microbiol* 2009, **73**:165-179.
  40. Uyar E, Kurokawa K, Yoshimura M, Ishikawa S, Ogasawara N, Oshima T: **Differential binding profiles of StpA in wild type and *hns* mutant cells: a comparative analysis of cooperative partners by chromatin immunoprecipitation-microarray analysis.** *J Bacteriol* 2009, **191**:2388-2391.
  41. Lucchini S, McDermott P, Thompson A, Hinton JC: **The H-NS-like protein StpA represses the RpoS (sigma 38) regulon during exponential growth of *Salmonella* Typhimurium.** *Mol Microbiol* 2009, **74**:1169-1186.
  42. Doyle M, Fookes M, Ivens A, Mangan MW, Wain J, Dorman CJ: **An H-NS-like stealth protein aids horizontal DNA transmission in bacteria.** *Science* 2007, **315**:251-252.
- How do A-T rich plasmids sneak under the radar and avoid titrating nucleoid proteins away from the host's chromosome? In this paper, the authors show that the "stealth" protein Sfh, which functions as a plasmid encoded H-NS homologue, plays a key role. Ref. [43] describes a genome-wide study of Sfh.
43. Dillon SC, Cameron ADS, Hokamp K, Lucchini S, Hinton JCD, Dorman CJ: **Genome-wide analysis of the H-NS and Sfh regulatory networks in *Salmonella* Typhimurium identifies a plasmid-encoded transcription silencing mechanism.** *Mol Microbiol* 2010, **76**:1250-1265.
  44. Stoebel DM, Free A, Dorman CJ: **Anti-silencing: overcoming H-NS-mediated repression of transcription in Gram-negative bacteria.** *Microbiology* 2008, **154**:2533-2545.
  45. Dorman CJ, Kane KA: **DNA bridging and antibridging: a role for bacterial nucleoid-associated proteins in regulating the expression of laterally acquired genes.** *FEMS Microbiol Rev* 2009, **33**:587-592.
  46. Basu D, Khare G, Singh S, Tyagi A, Khosla S, Mande SC: **A novel nucleoid-associated protein of *Mycobacterium tuberculosis* is a sequence homologue of GroEL.** *Nucleic Acids Res* 2009, **37**:4944-4954.
  47. Gordon BR, Imperial R, Wang L, Navarre WW, Liu J: **Lsr2 of *Mycobacterium* represents a novel class of H-NS-like proteins.** *J Bacteriol* 2009, **191**:2388-2391.
  48. Gordon BR, Li Y, Wang L, Sintsova A, van Bakel H, Tian S, Navarre WW, Xia B, Liu J: **Lsr2 is a nucleoid-associated protein that targets AT-rich sequences and virulence genes in *Mycobacterium tuberculosis*.** *Proc Natl Acad Sci* 2010, **107**:5154-5159.

49. Werlang IC, Schneider CZ, Mendonca JD, Palma MS, Basso LA, Santos OS: **Identification of Rv3852 as a nucleoid-associated protein in *Mycobacterium tuberculosis***. *Microbiology* 2009, **155**:2652-2663.
50. Sharadamma N, Harshavardhana Y, Singh P, Muniyappa K: ***Mycobacterium tuberculosis* nucleoid-associated DNA-binding protein H-NS binds with high-affinity to the Holliday junction and inhibits strand exchange promoted by RecA protein**. *Nucleic Acids Res* 2010, **38**:3555-3569.
51. Park HS, Ostberg Y, Johansson J, Wagner EG, Uhlin BE: **Novel role for a bacterial nucleoid protein in translation of mRNAs with suboptimal ribosome-binding sites**. *Genes Dev* 2010, **24**:1345-1350.
- This paper describes experiments to show that H-NS can act as a positive effector of gene expression by interacting with a specific messenger RNA and stimulating the initiation of translation. This observation may well resolve some longstanding anomalies in the literature concerning what H-NS does and does not do.
52. Corcoran CP, Dorman CJ: **DNA relaxation-dependent phase biasing of the *fim* genetic switch in *Escherichia coli* depends on the interplay of H-NS, IHF and LRP**. *Mol Microbiol* 2009, **74**:1071-1082.
53. Browning DF, Cole JA, Busby SJ: **Regulation by nucleoid-associated proteins at the *Escherichia coli nir* operon promoter**. *J Bacteriol* 2008, **190**:7258-7267.
54. Browning DF, Lee DJ, Spiro S, Busby SJ: **Down-regulation of the *Escherichia coli* K-12 *nrf* promoter by binding of the NsrR nitric oxide sensing transcription repressor to an upstream site**. *J Bacteriol* 2010, **192**:3824-3828.
55. Cho BK, Knight EM, Barrett CL, Palsson BO: **Genome-wide analysis of Fis binding in *Escherichia coli* indicates a causative role for A-/AT-tracts**. *Genome Res* 2008, **18**:900-910.
56. Stella S, Cascio D, Johnson RC: **The shape of the DNA minor groove directs binding by the DNA-bending protein Fis**. *Genes Dev* 2010, **24**:814-826.
- This paper presents a thorough structural analysis, which describes how DNA conformation, as well as DNA sequence, plays a key role in permitting Fis to interact with its DNA target.
57. Opel M, Aeling KA, Holmes WM, Johnson RC, Benham CJ, Hatfield GW: **Activation of transcription initiation from a stable RNA promoter by a Fis protein-mediated DNA structural transmission mechanism**. *Mol Microbiol* 2004, **53**:665-674.
58. Squire DJ, Xu M, Cole JA, Busby SJ, Browning DF: **Competition between NarL-dependent activation and Fis-dependent repression controls expression from the *Escherichia coli* *yeaR* and *ogt* promoter**. *Biochem J* 2009, **420**:249-257.
59. Grainger DC, Goldberg MD, Lee DJ, Busby SJ: **Selective repression by Fis and H-NS at the *Escherichia coli* *dps* promoter**. *Mol Microbiol* 2008, **68**:1366-1377.
60. Dame RT, Dorman CJ (Eds): *Bacterial Chromatin*. Springer; 2010.
61. Butala M, Busby SJ, Lee DJ: **DNA sampling: a method for probing protein binding at specific loci on bacterial chromosomes**. *Nucleic Acids Res* 2009, **37**:e37.